

DeltAg

Improving Plant Performance

THE ORIGINAL



Zinc Plus™

SOIL/FOLIAR

GUARANTEED ANALYSIS

0-0-0

Combined Sulfur (S) 4.90%
Soluble Zinc (Zn) 10.00%

Derived from Zinc Sulfate

Weight per gallon: 10.92 lb at 68 °F

Grams per liter: 1310 g at 20 °C

GENERAL INFORMATION: Zinc Plus is designed as a source of zinc to treat or prevent deficiencies for all field, row, vegetable, fruit, tree, vine, ornamental and turf crops.

CONTAINS NON-PLANT FOOD INGREDIENTS: Amino Acids, Enzymes, Vitamins and DeltAg Proprietary Complexing Agent.

Zinc Plus may be

- Foliar applied alone or with pesticides
- Used in drip irrigation systems
- Applied to soil via conventional methods

CAUTION: KEEP CONTAINER TIGHTLY CLOSED AND OUT OF REACH OF CHILDREN.

APPLICATION RATE:

Soil Incorporated: 32 to 64 ounces per acre.

Side-Dress: 16 to 32 ounces per acre.

Seed Treatment: 10 ounces per 100 pounds seed.

In-Furrow: 16 ounces per acre.

Foliar: 1 quart per acre

**Apply in a minimum 5 gallon volume for ample coverage.

* As much as 64 ounces per acre can be applied where deficiencies are observed.

Drip Irrigation: Apply at a rate of 0.25 ounce per 1,000 square feet of foliage weekly as needed.

Injection at 1 to 100: Add 2 ounces per gallon of mix tank water (156 ppm of product) weekly as needed.

Individual Trees & Plants:

Soil: Dilute 4 ounces in five gallons of water and inject into soil at least 10 inches deep in two to four locations under drip line.

Foliar: Dilute 4 ounces per each gallon of water and apply to foliage.

Application Timing: Field Crops: At planting or during seedling stages of crop development.

Orchards & Trees: Before leaf drop in Fall and again at Bud Break and repeat in three weeks.

COMPATIBILITY: This product is compatible with most commonly used pesticides. However, it is advisable to always check for compatibility.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Triple rinse container; empty rinsate into spray tank. Dispose of empty container according to approved local practices.

CONDITIONS OF SALE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material when not in strict accordance with directions.

SHAKE WELL BEFORE USING

DeltAg Formulations

P.O. Box 4812 Greenville, MS 38704

877 594 1500

www.deltagformulations.com

F1193

MR300-11